



Some Useful Techniques

Solutions

The Open University Press, Walton Hall, Milton Keynes.

First published 1975

Copyright © 1975. The Open University.

All rights reserved. No part of this work may be reproduced in any form, by mimeograph or any other means, without permission in writing from the publishers.

Produced in Great Britain by

Technical Filmsetters Europe Limited, 76 Great Bridgewater Street, Manchester M1 5JY.

ISBN 0 335 01257 4

This text forms part of the correspondence element of an Open University Third Level Course.

For general availability of supporting material referred to in this text, please write to the Director of Marketing, The Open University, P.O. Box 81, Milton Keynes, MK7 6AT.

Further information on Open University courses may be obtained from The Admissions Office, The Open University, P.O. Box 48, Milton Keynes, MK7 6AB.

Contents

	<i>Page</i>
0S.1 Ordinary Derivatives	4
0S.1.1 Differentiation	4
0S.1.2 Integration	9
0S.1.3 Differential Equations of First Order and Degree	12
0S.1.4 Differential Equations of First Order and Second Degree	14
0S.2 First-Order Partial Differentiation	15
0S.2.1 First-Order Partial Derivatives	15
0S.2.2 Integration of Partial Derivatives	17
0S.2.3 Partial Differential Operators	19
0S.2.4 Differentiation Along a Curve: the Chain Rule	20
0S.2.5 Derivatives “Following the Motion”	23
0S.2.6 Change of Coordinates	25
0S.3 Second-Order Partial Differentiation	30
0S.3.1 Second and Higher Order Partial Derivatives	30
0S.3.2 Solving Simple Second-Order Partial Differential Equations	31
0S.3.3 Transforming Second-Order Derivatives	34
0S.4 Vector Algebra and Line Integrals	40
0S.4.1 Vector Algebra	40
0S.4.2 Line Integrals	42
0S.5 Double and Volume Integrals	45
0S.5.1 Double Integrals	45
0S.5.2 Volume Integrals	47

0S.1 ORDINARY DERIVATIVES

0S.1.1 Differentiation

Solution to Exercise 1

- (i) $\sin' = \cos$ (see *Unit M100 12*) and so

$$\frac{d}{dx}(\sin x) = \boxed{\cos x}.$$

- (ii) The composite function rule is

$$(f \circ g)' = (f' \circ g) \times g'$$

(see *Unit M100 12*). Here $f = \exp$ and g is $x \mapsto (x - a)$, or just $(x - a)$ for short. Omitting the small circles as well, for brevity, we get

$$\begin{aligned} (\exp(x - a))' &= \exp'(x - a) \times (x - a)' \\ &= \exp(x - a) \times 1, \end{aligned}$$

since $\exp' = \exp$ (see *Unit M100 12*),

$$= \exp(x - a).$$

We could alternatively have written the composite function rule in Leibniz notation,

$$\frac{dz}{dx} = \frac{dz}{dy} \frac{dy}{dx},$$

(often called the *chain rule* for functions of one variable), where $z(y) = \exp y$ and $y(x) = x - a$. Then

$$\begin{aligned} \frac{dz}{dx} &= \exp y \times 1 \\ &= \exp(x - a). \end{aligned}$$

In either case we get

$$\frac{d}{dx}(\exp(x - a)) = \boxed{\exp(x - a)}.$$

- (iii) We can use either of the methods in part (ii) above. Using the Leibniz chain rule with $z(y) = \sqrt{y}$ and $y(x) = \sin x$,

$$\frac{dz}{dx} = \frac{1}{2}y^{-\frac{1}{2}} \times \cos x,$$

so that

$$\frac{d}{dx}(\sqrt{\sin x}) = \boxed{\frac{\cos x}{2\sqrt{\sin x}}}.$$

- (iv) The definitions we need are

$$\cosh x = \frac{1}{2}(e^x + e^{-x}), \quad \sinh x = \frac{1}{2}(e^x - e^{-x}),$$

from which we see that the derived function of \cosh is \sinh (*not* minus \sinh ; the analogy between trigonometric and hyperbolic functions is imperfect). Applying the chain rule as above with $z(y) = \cosh y$, $y(x) = x^2$,

$$\frac{dz}{dx} = \sinh y \times 2x,$$

so that

$$\frac{d}{dx}(\cosh x^2) = \boxed{2x \sinh x^2}.$$

Solution to Exercise 2

Using the chain rule with $z(y) = 1/y$, $y(x) = \cos x$,

$$\begin{aligned}\frac{dz}{dx} &= -\frac{1}{y^2} \times (-\sin x), \\ &= \frac{\sin x}{\cos^2 x}\end{aligned}$$

so that

$$\frac{d}{dx}(\sec x) = \boxed{\tan x \sec x}.$$

If $h = f/g$, then

$$h' = \frac{f'g - fg'}{g^2}.$$

This is the form of the quotient rule stated in *Unit M100 12*. In Leibniz notation, if $u = u(x)$ and $v = v(x)$ then

$$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}.$$

Putting $u(x) = \sin x$, $v(x) = \cos x$,

$$\begin{aligned}\frac{d}{dx}\left(\frac{\sin x}{\cos x}\right) &= \frac{\cos x(\cos x) - \sin x(-\sin x)}{\cos^2 x} \\ &= \frac{1}{\cos^2 x},\end{aligned}$$

so that

$$\frac{d}{dx}(\tan x) = \boxed{\sec^2 x}.$$

This could also be written in other equivalent forms, such as $1 + \tan^2 x$.

Solution to Exercise 3

Since

$$x = \frac{1}{2}at^2,$$

we have

$$\begin{aligned}\frac{dx}{dt} &= at \\ &= \sqrt{2ax},\end{aligned}$$

so that

$$v(t) = \boxed{at},$$

$$v(x) = \boxed{\sqrt{2ax}}.$$

If $t = 4$, then $v = 4a$, so that

$$v(t = 4) = \boxed{4a}.$$

If $x = 4$, then $v = \sqrt{8a}$, so that

$$v(x = 4) = \boxed{\sqrt{8a}}.$$

$v(t = x)$ is the expression for $v(t)$ with x replacing t ,

$$v(t = x) = \boxed{ax}.$$

$v(x = t)$ is the expression for $v(x)$ with t replacing x ,

$$v(x = t) = \boxed{\sqrt{2at}}.$$

$v|_{x=3}$ is the same as $v(x = 3)$, that is

$$v|_{x=3} = \boxed{\sqrt{6a}}.$$

The derivative of v with respect to t is $d(at)/dt$, so that

$$\frac{dv}{dt} = \boxed{a}.$$

The derivative of v with respect to x is $d(\sqrt{2ax})/dx$, which is

$$\sqrt{2a} \frac{d}{dx}(x^{\frac{1}{2}}) = \frac{1}{2} \sqrt{2a} x^{-\frac{1}{2}},$$

or

$$\frac{dv}{dx} = \boxed{\sqrt{\frac{a}{2x}}}.$$

If $x = 1$, then $dv/dx = \sqrt{(a/2)}$, or

$$\left. \frac{dv}{dx} \right|_{x=1} = \boxed{\sqrt{\frac{a}{2}}}.$$

If $t = t_0$, then $x = \frac{1}{2}at_0^2$, so that

$$\begin{aligned} \left. \frac{dv}{dx} \right|_{t=t_0} &= \left. \frac{dv}{dx} \right|_{x=\frac{1}{2}at_0^2} \\ &= \sqrt{\frac{a}{2(\frac{1}{2}at_0^2)}}, \end{aligned}$$

giving

$$\left. \frac{dv}{dx} \right|_{t=t_0} = \boxed{\frac{1}{t_0}}.$$

Solution to Exercise 4

From above,

$$\frac{dx}{dt} = \boxed{3t^2}.$$

We have $v(t) = 3t^2$, $v(x) = 3x^{\frac{2}{3}}$, so that

$$\frac{dv}{dt} = \boxed{6t},$$

$$\frac{dv}{dx} = \boxed{2x^{-\frac{1}{3}}}.$$

Expressing dv/dx in terms of t ,

$$\frac{dv}{dx} = \frac{2}{t},$$

giving

$$\frac{dv}{dx} \frac{dx}{dt} = \boxed{6t}.$$

$$\text{Is } \frac{dv}{dt} = \frac{dv}{dx} \frac{dx}{dt} ? \quad \boxed{\text{Yes}}$$

Solution to Exercise 5

Since $x = \frac{1}{2}at^2$ in Exercise 3,

$$x(t) \text{ is } t \mapsto \boxed{\frac{1}{2}at^2}.$$

Differentiating with respect to t ,

$$\frac{dx}{dt} = \boxed{at}.$$

From Exercise 3,

$$\frac{dv}{dt} = \boxed{a},$$

and

$$\frac{dv}{dx} = \boxed{\sqrt{\frac{a}{2x}}}.$$

Expressing this in terms of t ,

$$\frac{dv}{dx} = \frac{1}{t},$$

so that

$$\frac{dv}{dx} \frac{dx}{dt} = \boxed{a}.$$

$$\text{Is } \frac{dv}{dt} = \frac{dv}{dx} \frac{dx}{dt} ? \quad \boxed{\text{Yes}}$$

Solution to Exercise 6

If $z = z(y)$, $y = y(x)$ and $x = x(t)$, we know that the chain rule gives both

$$\frac{dz}{dt} = \frac{dz}{dy} \frac{dy}{dt}$$

and

$$\frac{dy}{dt} = \frac{dy}{dx} \frac{dx}{dt}.$$

Combining these two,

$$\frac{dz}{dt} = \boxed{\frac{dz}{dy} \frac{dy}{dx} \frac{dx}{dt}}.$$

Put $z = \ln y$, $y = \cos x$, $x = e^{2t}$. Then

$$\frac{dz}{dy} = \frac{1}{y} = \frac{1}{\cos x} = \frac{1}{\cos e^{2t}},$$

$$\frac{dy}{dx} = -\sin x = -\sin e^{2t},$$

$$\frac{dx}{dt} = 2e^{2t},$$

so that

$$\begin{aligned} \frac{dz}{dt} &= \frac{1}{\cos e^{2t}} (-\sin e^{2t}) 2e^{2t} \\ &= -2e^{2t} \tan e^{2t}. \end{aligned}$$

Thus,

$$\frac{d}{dt}(\ln \cos e^{2t}) = \boxed{-2e^{2t} \tan e^{2t}}.$$

Solution to Exercise 7

If $y(x) = \arctan x$, then

$$x(y) = \boxed{\tan y};$$

hence, by Exercise 2,

$$\frac{dx}{dy}(y) = \boxed{\sec^2 y};$$

hence, by the chain rule,

$$\frac{dy}{dx}(y) = \boxed{\cos^2 y}.$$

Since $\sec^2 y = 1 + \tan^2 y$,

$$\cos^2 y = \frac{1}{1 + x^2};$$

thus

$$\frac{dy}{dx}(x) = \boxed{\frac{1}{1 + x^2}},$$

so that

$$\frac{d}{dx}(\arctan x) = \boxed{\frac{1}{1+x^2}}.$$

0S.1.2 Integration

Solution to Exercise

The methods described below are all based very firmly on the M201 supplementary handbook, *Techniques of Integration (TI)*. If you know a quicker method, so much the better. The constant of integration is omitted in each indefinite integral.

- (i) $\int x \, dx$. The integrand x is a polynomial function of x , so we can use the formula in section III.1 on page 8 of *TI*, with $a_1 = 1$, and $a_i = 0$ for all $i \neq 1$, to get

$$\int x \, dx = \boxed{\frac{1}{2}x^2}.$$

- (ii) $\int t^2 \, dt$. The integrand is a polynomial function of t . We use the same formula as in part (i), but on this occasion t replaces x and we have $a_2 = 1$ and $a_i = 0$ for all $i \neq 2$. Thus

$$\int t^2 \, dt = \boxed{\frac{1}{3}t^3}.$$

- (iii) $\int e^{-x} \, dx$. The integrand involves an exponential function, so we look at section III.5 on page 14 of *TI*. Formula 5.2(i) with $a = -1$, $b = 0$ gives

$$\int e^{-x} \, dx = \boxed{-e^{-x}}.$$

Alternatively formula 5.2(iii) with $a = -1$, $n = 0$ (or even formula 5.2(vi) with $a = e^{-1}$) gives the same answer.

- (iv) $\int \frac{1}{1+x^2} \, dx$. The integrand is a rational function of x , so we can use section III.2 on page 8 of *TI*. The first relevant paragraph is 2.3, with $p = 0$, $q = 1$, $b = 0$, $c = 1$ (and note that $b^2 < 4c$; if this had not been so we would have had to go on to paragraph 2.4). The recommended substitution $t = x + \frac{1}{2}b$ is just $t = x$, i.e. we can proceed to the next step immediately. Using the second of the two given standard integrals, with $c = 1$ and x in place of t , we get

$$\int \frac{1}{1+x^2} \, dx = \boxed{\arctan x}.$$

The arctan function was mentioned in Exercise 7 of subsection 0.1.1, and you may have been able to answer this question by using the result of that exercise. A graph of the arctan function is shown on page 7 of *TI*.

- (v) $\int \frac{1}{4+x^2} \, dx$. As in part (iv), we are led to paragraph 2.3 in part III of *TI*. This time we have $p = 0$, $q = 1$, $b = 0$, $c = 4$, and so the second standard integral at the top of page 9 gives

$$\int \frac{1}{4+x^2} \, dx = \boxed{\frac{1}{2} \arctan \frac{1}{2}x}.$$

- (vi) $\int \sin \alpha x \sin \beta x dx$. The integrand involves trigonometric functions, so we turn to section III.5 on page 14 of *TI*. Using the first formula in paragraph 5.1, with α, β replacing a, b respectively, brings the integral to the form

$$\frac{1}{2} \int \cos(\alpha - \beta)x dx - \frac{1}{2} \int \cos(\alpha + \beta)x dx.$$

The next relevant formula is 5.2(i) with $a = 0$ and b put equal to $\alpha - \beta$ and $\alpha + \beta$ in turn. This gives

$$\int \sin \alpha x \sin \beta x dx = \boxed{\frac{1}{2} \left[\frac{\sin(\alpha - \beta)x}{\alpha - \beta} - \frac{\sin(\alpha + \beta)x}{\alpha + \beta} \right]}.$$

- (vii) $\int_0^\pi \sin mx \sin nx dx$. We have two cases to consider, depending upon whether or not m and n are equal. Suppose $m \neq n$. We then have from the answer to part (vi), with m, n replacing α, β respectively,

$$\begin{aligned} \int_0^\pi \sin mx \sin nx dx &= \frac{1}{2} \left[\frac{\sin(m - n)x}{m - n} - \frac{\sin(m + n)x}{m + n} \right]_0^\pi \\ &= 0, \end{aligned}$$

since the sine of any integer multiple of π is zero. Suppose next that $m = n$. Our integrand is now $\sin^2 nx$, and using the first formula in paragraph 5.1 (section III.5 on page 14 of *TI*) with $a = b = n$,

$$\begin{aligned} \int_0^\pi \sin^2 nx dx &= \frac{1}{2} \int_0^\pi (1 - \cos 2nx) dx \\ &= \frac{1}{2} \left[x - \frac{1}{2n} \sin 2nx \right]_0^\pi, \end{aligned}$$

having used formula 5.2(i) with $a = 0, b = 2n$ to integrate the second term of the integrand. Thus

$$\int_0^\pi \sin^2 nx dx = \frac{1}{2}\pi,$$

so that

$$\int_0^\pi \sin mx \sin nx dx = \begin{cases} \boxed{\frac{1}{2}\pi} & \text{if } m = n, \\ \boxed{0} & \text{if } m \neq n. \end{cases}$$

- (viii) $\int_0^\pi \cos x \sin nx dx$. We again start with paragraph 5.1 of section III.5 on page 14 of *TI*. Using the third formula in this paragraph, with $a = n, b = 1$, our integral becomes

$$\frac{1}{2} \int_0^\pi \sin(n - 1)x dx + \frac{1}{2} \int_0^\pi \sin(n + 1)x dx.$$

We then use formula 5.2(ii) with $a = 0$ and b equal to $n - 1, n + 1$ in turn to get

$$\int_0^\pi \cos x \sin nx dx = \frac{1}{2} \left[-\frac{\cos(n - 1)x}{n - 1} - \frac{\cos(n + 1)x}{n + 1} \right]_0^\pi$$

provided that $n \neq 1$ (if $n = 1$, we see that the first integrand above is zero, so that the first term in the square bracket should be replaced by zero in this case). Since $\cos m\pi = (-1)^m$ if m is an integer,

$$\int_0^\pi \cos x \sin nx dx = \begin{cases} \boxed{\frac{1}{n - 1} + \frac{1}{n + 1}} & \text{if } n \text{ is even,} \\ \boxed{0} & \text{if } n \text{ is odd.} \end{cases}$$

- (ix) $\int_0^\pi x \sin nx \, dx$. We can perform the integration by using formula 5.2(v) of section III.5 in *TI* with n replaced by 1 and a replaced by n . This gives

$$\begin{aligned}\int_0^\pi x \sin nx \, dx &= \left[-\frac{x}{n} \cos nx + \frac{1}{n^2} \sin nx \right]_0^\pi \\ &= -\frac{\pi}{n} \cos n\pi,\end{aligned}$$

so that

$$\int_0^\pi x \sin nx \, dx = \boxed{(-1)^{n+1} \frac{\pi}{n}}.$$

- (x) $\int_{-\pi}^\pi x^2 \cos nx \, dx$. Using formula 5.2(iv) of section III.5 with n replaced by 2 and a replaced by n ,

$$\begin{aligned}\int_{-\pi}^\pi x^2 \cos nx \, dx &= \left[\left(\frac{x^2}{n} - \frac{2}{n^3} \right) \sin nx + \frac{2x}{n^2} \cos nx \right]_{-\pi}^\pi \\ &= \frac{2\pi}{n^2} \cos n\pi - \frac{2(-\pi)}{n^2} \cos(-n\pi) \\ &= \frac{4\pi}{n^2} \cos n\pi,\end{aligned}$$

so that

$$\int_{-\pi}^\pi x^2 \cos nx \, dx = \boxed{4(-1)^n \frac{\pi}{n^2}}.$$

- (xi) $\int_{-\pi}^\pi f(x) \sin nx \, dx$. Since $f(x) = 0$ for $-\pi \leq x \leq 0$ and $f(x) = 1$ for $0 < x \leq \pi$, we break our integral up into two parts:

$$\begin{aligned}\int_{-\pi}^\pi f(x) \sin nx \, dx &= \int_{-\pi}^0 0 \times \sin nx \, dx + \int_0^\pi 1 \times \sin nx \, dx \\ &= \int_0^\pi \sin nx \, dx \\ &= \left[-\frac{1}{n} \cos nx \right]_0^\pi,\end{aligned}$$

having used formula 5.2(ii) with $a = 0, b = n$. Since

$$\left[-\frac{1}{n} \cos nx \right]_0^\pi = -\frac{1}{n} \cos n\pi + \frac{1}{n},$$

it follows that

$$\int_{-\pi}^\pi f(x) \sin nx \, dx = \boxed{\frac{1}{n}(1 - (-1)^n)}.$$

- (xii) $\int_0^1 f(x) \sin n\pi x \, dx$. Since $f(x) = x$ for $0 \leq x \leq \frac{1}{2}$ and $f(x) = 1 - x$ for $\frac{1}{2} \leq x \leq 1$, we break our integral into two parts:

$$\begin{aligned}\int_0^1 f(x) \sin n\pi x \, dx &= \int_0^{\frac{1}{2}} f(x) \sin n\pi x \, dx + \int_{\frac{1}{2}}^1 f(x) \sin n\pi x \, dx \\ &= \int_0^{\frac{1}{2}} x \sin n\pi x \, dx + \int_{\frac{1}{2}}^1 (1 - x) \sin n\pi x \, dx.\end{aligned}$$

We see by putting $n = 1$, $a = n\pi$ in formula 5.2(v) in section III.5 on page 14 of **TI** that

$$\int x \sin n\pi x \, dx = -\frac{x}{n\pi} \cos n\pi x + \frac{1}{n^2\pi^2} \sin n\pi x.$$

By putting $n = 0$, $a = n\pi$ in the same formula,

$$\int \sin n\pi x \, dx = -\frac{1}{n\pi} \cos n\pi x.$$

Thus,

$$\begin{aligned} \int_0^1 f(x) \sin n\pi x \, dx &= \left[-\frac{x}{n\pi} \cos n\pi x + \frac{1}{n^2\pi^2} \sin n\pi x \right]_0^{\frac{1}{2}} \\ &\quad + \left[-\frac{1}{n\pi} \cos n\pi x + \frac{x}{n\pi} \cos n\pi x - \frac{1}{n^2\pi^2} \sin n\pi x \right]_{\frac{1}{2}}^1 \\ &= -\frac{1}{2n\pi} \cos \frac{1}{2}n\pi + \frac{1}{n^2\pi^2} \sin \frac{1}{2}n\pi \\ &\quad + \frac{1}{2n\pi} \cos \frac{1}{2}n\pi + \frac{1}{n^2\pi^2} \sin \frac{1}{2}n\pi \\ &= \frac{2}{n^2\pi^2} \sin \frac{1}{2}n\pi. \end{aligned}$$

Now $\sin \frac{1}{2}n\pi$ is zero if n is even. If n is odd, $\sin \frac{1}{2}n\pi = 1$ if $\frac{1}{2}(n-1)$ is even and $\sin \frac{1}{2}n\pi = -1$ if $\frac{1}{2}(n-1)$ is odd.

Therefore,

$$\int_0^1 f(x) \sin n\pi x \, dx = \begin{cases} 0 & \text{if } n \text{ is even,} \\ (-1)^{\frac{1}{2}(n-1)} \frac{2}{n^2\pi^2} & \text{if } n \text{ is odd.} \end{cases}$$

0S.1.3 Differential Equations of First Order and Degree

Solution to Exercise

- (i) Integrating both sides we obtain

$$x(t) = \boxed{\frac{1}{3}t^3 + A}.$$

- (ii) Separating variables (see page 20 of *Unit M100 24* or section 4.2 of **SS** if necessary),

$$\int e^{-x} \, dx = \int dt + A,$$

or

$$-e^{-x} = t + A.$$

Therefore

$$x(t) = \boxed{-\ln(-t - A)} \quad t \in (-\infty, -A).$$

Notice that in this case no solution has as domain the whole of R . This commonly happens with nonlinear differential equations, for which the simple uniqueness theorem studied in *Units M201 4* and *M201 33* may not apply.

(iii) Separating variables,

$$\int x \, dx = \int t \, dt + C,$$

giving

$$\frac{1}{2}x^2 = \frac{1}{2}t^2 + C,$$

so that

$$x(t) = \boxed{\pm \sqrt{t^2 + A}},$$

where $A = 2C$, and $t \in R$ if $A \geq 0$, $t \in (\sqrt{-A}, \infty) \cup (-\infty, -\sqrt{-A})$ if $A < 0$.

(iv) Separating variables,

$$\int \frac{1}{1+x^2} dx = \int dt + A,$$

and using part (iv) of the Exercise in subsection 0.1.2 this gives

$$\arctan x = t + A,$$

or

$$x(t) = \boxed{\tan(t + A)} \quad t \in (-A - \frac{1}{2}\pi, -A + \frac{1}{2}\pi).$$

(v) Separating variables,

$$\int \frac{1}{4+x^2} dx = \int dt + C,$$

so that using part (v) of the Exercise in subsection 0.1.2 gives

$$\frac{1}{2}\arctan \frac{1}{2}x = t + C,$$

or

$$x(t) = \boxed{2 \tan(2t + A)} \quad t \in (-\frac{1}{2}A - \frac{1}{4}\pi, -\frac{1}{2}A + \frac{1}{4}\pi),$$

where $A = 2C$.

(vi) The equation can be written as

$$\frac{dx}{dt} - \frac{1}{t}x = 0,$$

so that an integrating factor is

$$\exp \int \left(-\frac{1}{t}\right) dt = \exp(-\ln t) = \frac{1}{t}$$

(see lines 6 to 10 on page 97 of *K* if necessary). Multiplying through by this factor, we see that the equation can be put in the form

$$\frac{d}{dt} \left(\frac{x}{t} \right) = 0,$$

which has solution

$$x(t) = \boxed{At}.$$

0S.1.4 Differential Equations of First Order and Second Degree

Solution to Exercise

- (i) Factorizing the equation gives

$$\left(2\frac{dx}{dt} - 1\right)\left(\frac{dx}{dt} + 1\right) = 0,$$

so that

$$\frac{dx}{dt} = \frac{1}{2} \text{ or } -1.$$

(Alternatively you can use the general quadratic equation formula

$$\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

to achieve this result.)

Integrating,

$$x(t) = \boxed{-t + A} \quad \text{or} \quad \boxed{\frac{1}{2}t + A}.$$

- (ii) Factorizing gives

$$\left(\frac{dx}{dt} - e^x\right)\left(\frac{dx}{dt} - t^2\right) = 0,$$

so that

$$\frac{dx}{dt} = e^x \quad \text{or} \quad t^2.$$

Hence from parts (i) and (ii) of the Exercise in subsection 0.1.3,

$$x(t) = \boxed{-\ln(-t - A)} \quad \text{or} \quad \boxed{\frac{1}{3}t^3 + A},$$

with the restriction that $t \in (-\infty, -A)$ in the first case.

0S.2 FIRST-ORDER PARTIAL DIFFERENTIATION

0S.2.1 First-Order Partial Derivatives

Solution to Exercise 1

$$\begin{aligned}
 \text{(i)} \quad \frac{\partial u}{\partial x}(x, y) &= \boxed{2x} \\
 \frac{\partial u}{\partial y}(x, y) &= \boxed{2y} \\
 \text{(ii)} \quad \frac{\partial u}{\partial x}(x, y) &= \boxed{y^2 e^{xy}} \\
 \frac{\partial u}{\partial y}(x, y) &= \boxed{(1 + xy)e^{xy}} \\
 \text{(iii)} \quad \frac{\partial u}{\partial x}(x, y) &= \boxed{\cos(x - y)} \\
 \frac{\partial u}{\partial y}(x, y) &= \boxed{-\cos(x - y)} \\
 \text{(iv)} \quad \frac{\partial u}{\partial x}(x, y) &= \boxed{f'(x - y)} \\
 \frac{\partial u}{\partial y}(x, y) &= \boxed{-f'(x - y)}
 \end{aligned}$$

Solution to Exercise 2

$$\begin{aligned}
 \frac{\partial}{\partial x}[p(x + ct) + q(x - ct)] &= \boxed{p'(x + ct) + q'(x - ct)}, \\
 \frac{\partial}{\partial t}[p(x + ct) + q(x - ct)] &= \boxed{cp'(x + ct) - cq'(x - ct)}.
 \end{aligned}$$

Solution to Exercise 3

- (i) Suppose that G is a primitive of g , so that by the fundamental theorem of calculus, $G' = g$. Then

$$\begin{aligned}
 \int_0^{x+ct} g(\bar{x}) d\bar{x} &= \left[G(\bar{x}) \right]_0^{x+ct} \\
 &= G(x + ct) - G(0),
 \end{aligned}$$

so that

$$\begin{aligned}
 \frac{\partial}{\partial x} \int_0^{x+ct} g(\bar{x}) d\bar{x} &= \frac{\partial}{\partial x} [G(x + ct) - G(0)] \\
 &= G'(x + ct),
 \end{aligned}$$

or

$$\frac{\partial}{\partial x} \int_0^{x+ct} g(\bar{x}) d\bar{x} = \boxed{g(x + ct)}.$$

- (ii) This is the same as for part (i), with $x - ct$ replacing $x + ct$ throughout. Thus

$$\frac{\partial}{\partial x} \int_0^{x-ct} g(\bar{x}) d\bar{x} = \boxed{g(x - ct)}.$$

(iii) Since

$$\int_{x-ct}^{x+ct} g(\bar{x}) d\bar{x} = \int_0^{x+ct} g(\bar{x}) d\bar{x} - \int_0^{x-ct} g(\bar{x}) d\bar{x},$$

we have from parts (i) and (ii)

$$\frac{\partial}{\partial x} \int_{x-ct}^{x+ct} g(\bar{x}) d\bar{x} = \boxed{g(x+ct) - g(x-ct)}.$$

(iv) Defining G as before,

$$\frac{\partial}{\partial t} \int_{x-ct}^{x+ct} g(\bar{x}) d\bar{x} = \frac{\partial}{\partial t} [G(x+ct) - G(x-ct)].$$

Using the second result of Exercise 2 with $p = G$, $q = -G$,

$$\frac{\partial}{\partial t} \int_{x-ct}^{x+ct} g(\bar{x}) d\bar{x} = \boxed{cg(x+ct) + cg(x-ct)}.$$

Solution to Exercise 4

If $u(x, t) = \sin(x^2 - t)$, put $t = 0$ to get $u(x, 0) = u|_{t=0}$:

$$u|_{t=0} = \boxed{\sin x^2}.$$

The partial derivative of u with respect to t is

$$\frac{\partial u}{\partial t}(x, t) = -\cos(x^2 - t),$$

so that

$$\left. \frac{\partial u}{\partial t} \right|_{t=0} = \boxed{-\cos x^2}.$$

The partial derivative of u with respect to x is

$$\frac{\partial u}{\partial x}(x, t) = \boxed{2x \cos(x^2 - t)}.$$

Also,

$$u|_{x=0} = \boxed{-\sin t},$$

$$\left. \frac{\partial u}{\partial x} \right|_{x=1} = \boxed{2 \cos(1 - t)}.$$

0S.2.2 Integration of Partial Derivatives

Solution to Exercise 1

- (i) Since partial differentiation of a function of x and y with respect to x is performed by treating y as a constant,

$$\frac{\partial u}{\partial x} = \boxed{0}.$$

- (ii) As in part (i),

$$\frac{\partial u}{\partial x} = \boxed{0}.$$

- (iii) As in parts (i) and (ii),

$$\frac{\partial u}{\partial x} = \boxed{0}.$$

Solution to Exercise 2

The general solution is the function we differentiated in Exercise 1(iii):

$$u(x, y) = \boxed{\begin{array}{l} f(y), \\ \text{where } f \text{ is an arbitrary function} \\ \text{of one variable.} \end{array}}$$

Solution to Exercise 3

The general solution is

$$u(x, y) = \boxed{\begin{array}{l} f(x), \\ \text{where } f \text{ is an arbitrary function} \\ \text{of one variable.} \end{array}}$$

A brief justification:

For fixed x , consider

$$y \mapsto u(x, y).$$

This has derived function

$$y \mapsto \frac{\partial u}{\partial y}(x, y),$$

which is

$$y \mapsto 0$$

by the differential equation. Hence $y \mapsto u(x, y)$ is a constant function (regarding x as a fixed number) so that

$$\begin{aligned} u(x, y) &= u(x, 0) \\ &= f(x) \quad (\text{say}). \end{aligned}$$

Solution to Exercise 4

- (i) The general solution of the associated homogeneous equation is

$$u(x, y) = f(y)$$

for arbitrary f . If we regard y as fixed, a particular primitive of the function

$$x \mapsto y$$

is

$$x \mapsto xy,$$

so that the general solution we require is

$$u(x, y) = \boxed{xy + f(y)}.$$

- (ii) Since for fixed
- x
- the function

$$y \mapsto xy$$

has a primitive

$$y \mapsto \frac{1}{2}xy^2,$$

the general solution is

$$u(x, y) = \boxed{\frac{1}{2}xy^2 + f(x)}.$$

- (iii) Since for fixed
- t
- the function

$$x \mapsto x^2 + t^2$$

has a primitive

$$x \mapsto \frac{1}{3}x^3 + xt^2,$$

the general solution is

$$u(x, t) = \boxed{\frac{1}{3}x^3 + xt^2 + f(t)}.$$

- (iv) Since for fixed
- ξ
- the function

$$\eta \mapsto r(\eta)$$

has a primitive

$$\eta \mapsto \int r(\eta) d\eta,$$

the general solution is

$$w(\xi, \eta) = \boxed{\int r(\eta) d\eta + f(\xi)}.$$

Solution to Exercise 5

- (i) By the standard formula (see lines 6 to 10 on page 97 of
- K
-), an integrating factor for the ordinary differential equation is

$$\exp \int \left(-\frac{1}{x} \right) dx = \exp(-\ln x),$$

so that the integrating factor is

$$\boxed{\frac{1}{x}}.$$

(ii) Using this factor for the partial differential equation we obtain

$$\frac{1}{x} \frac{\partial u}{\partial x} - \frac{u}{x^2} = 0,$$

or

$$\frac{\partial}{\partial x} \left(\frac{u(x, y)}{x} \right) = 0.$$

Thus

$$\frac{u(x, y)}{x} = f(y),$$

and the general solution is

$$u(x, y) = \boxed{xf(y)}.$$

Solution to Exercise 6

(i) The integrating factor is

$$\exp \int \left(-\frac{1}{y} \right) dx = \boxed{\exp \left(-\frac{x}{y} \right)}.$$

(ii) The partial differential equation becomes

$$\frac{\partial}{\partial x} \left(u(x, y) \exp \left(-\frac{x}{y} \right) \right) = 0,$$

so that the general solution is

$$u(x, y) = \boxed{e^{x/y} f(y)}.$$

0S.2.3 Partial Differential Operators

Solution to Exercise 1

Since the operator is to be first-order, we are limited to first- and zeroth-order derivatives, and there can be no products of the first-order derivatives since this would contravene the linearity of L . Thus the most general form possible for L is

$$L = \boxed{a(x, y) \frac{\partial}{\partial x} + b(x, y) \frac{\partial}{\partial y} + c(x, y)},$$

for arbitrary functions of two variables a , b and c .

Solution to Exercise 2

(i) Since

$$\frac{\partial}{\partial x} (x + y)^2 = \frac{\partial}{\partial y} (x + y)^2 = 2(x + y),$$

it follows that

$$\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} \right) (x + y)^2 = \boxed{4(x + y)}.$$

(ii) Since

$$\frac{\partial}{\partial x}(x - y)^2 = -\frac{\partial}{\partial y}(x - y)^2 = 2(x - y),$$

we have

$$\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y}\right)(x - y)^2 = \boxed{0}.$$

(iii) Since

$$\frac{\partial}{\partial x} \sin(x - y) = -\frac{\partial}{\partial y} \sin(x - y) = \cos(x - y),$$

we have

$$\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y}\right) \sin(x - y) = \boxed{0}.$$

(iv) Since

$$\frac{\partial}{\partial x} F(x, y) = -\frac{\partial}{\partial y} F(x - y) = F'(x - y),$$

we have

$$\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y}\right) F(x - y) = \boxed{0}.$$

0S.2.4 Differentiation Along a Curve: the Chain Rule

Solution to Exercise 1

(i) Put

$$x = 2t, y = \sin t$$

into the potential energy $\phi = x^2 + y^3$ to get

$$\phi(t) = \boxed{4t^2 + \sin^3 t}.$$

(ii) Differentiating this expression for $\phi(t)$ gives

$$\phi'(t) = \frac{d\phi}{dt} = \boxed{8t + 3 \sin^2 t \cos t}.$$

Solution to Exercise 2

(i) We have

$$z = x^2 + y^3,$$

and

$$x = \xi(t), y = \eta(t),$$

$$\text{so that } z(t) = \boxed{[\xi(t)]^2 + [\eta(t)]^3}.$$

(ii) Differentiating with respect to t gives

$$z'(t) = \frac{dz}{dt} = \boxed{2\xi(t)\xi'(t) + 3[\eta(t)]^2\eta'(t)}$$

Solution to Exercise 3

- (i) Since
- $x = \xi(t)$
- , we have

$$\frac{dx}{dt} = \boxed{\xi'(t)}.$$

- (ii) If
- $z = x^2 + y^3$
- , we have

$$\frac{\partial z}{\partial x} = \boxed{2x}.$$

- (iii) We want
- $\partial z/\partial x$
- evaluated at
- $x = \xi(t)$
- ,
- $y = \eta(t)$
- . From part (ii) we have

$$\frac{\partial z}{\partial x}(\xi(t), \eta(t)) = \boxed{2\xi(t)}.$$

- (iv) By proceeding as in parts (i), (ii) and (iii) we find that

$$\frac{dy}{dt} = \eta'(t),$$

$$\frac{\partial z}{\partial y}(\xi(t), \eta(t)) = 3[\eta(t)]^2,$$

so that at $(\xi(t), \eta(t))$ on the surface,

$$\frac{\partial z}{\partial x} \frac{dx}{dt} + \frac{\partial z}{\partial y} \frac{dy}{dt} = \boxed{2\xi(t)\xi'(t) + 3[\eta(t)]^2\eta'(t)}.$$

This is the same as the formula for dz/dt in Exercise 2(ii).

Solution to Exercise 4

- (i) This is the case in Exercises 2 and 3 (if
- z
- is replaced by
- u
-). Since the solutions to Exercises 2(ii) and 3(iv) are identical, we see that the boxed formula holds in this case.

Yes

- (ii) The result is also true in general; a demonstration is given in Appendix I of
- Unit MST 282 7*
- on pages 34, 35.

Yes

Solution to Exercise 5

The analogue of the formula in Exercise 4 for the case where u is a function of three variables x, y, z is

$$\boxed{\frac{du}{dt} = \frac{\partial u}{\partial x} \frac{dx}{dt} + \frac{\partial u}{\partial y} \frac{dy}{dt} + \frac{\partial u}{\partial z} \frac{dz}{dt}}$$

Solution to Exercise 6

(i) We have

$$\frac{\partial u}{\partial x} = 2x = 2t^2, \quad \frac{dx}{dt} = 2t,$$

$$\frac{\partial u}{\partial y} = 2y = 2(t-1) \quad \frac{dy}{dt} = 1,$$

$$\frac{\partial u}{\partial z} = 2z = 2t^3, \quad \frac{dz}{dt} = 3t^2,$$

so that

$$\frac{\partial u}{\partial x} \frac{dx}{dt} + \frac{\partial u}{\partial y} \frac{dy}{dt} + \frac{\partial u}{\partial z} \frac{dz}{dt} = (2t^2)(2t) + 2(t-1) + (2t^3)(3t^2),$$

giving

$$\frac{du}{dt} = \boxed{6t^5 + 4t^3 + 2t - 2}.$$

(ii) Writing

$$u = (t^2)^2 + (t-1)^2 + (t^3)^2$$

or

$$u(t) = \boxed{t^6 + t^4 + t^2 - 2t + 1},$$

we have

$$\frac{du}{dt} = \boxed{6t^5 + 4t^3 + 2t - 2}.$$

The two methods agree.

Solution to Exercise 7

(i) By definition we have

$$\text{grad } u = \frac{\partial u}{\partial x} \mathbf{i} + \frac{\partial u}{\partial y} \mathbf{j} + \frac{\partial u}{\partial z} \mathbf{k},$$

so that by reference to the solution of Exercise 6,

$$\text{grad } u = \boxed{2(t^2 \mathbf{i} + (t-1) \mathbf{j} + t^3 \mathbf{k})}.$$

(ii) If $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$, then

$$\frac{d\mathbf{r}}{dt} = \frac{dx}{dt} \mathbf{i} + \frac{dy}{dt} \mathbf{j} + \frac{dz}{dt} \mathbf{k},$$

or

$$\frac{d\mathbf{r}}{dt} = \boxed{2t \mathbf{i} + \mathbf{j} + 3t^2 \mathbf{k}}.$$

$$\begin{aligned} \text{(iii)} \quad \text{grad } u \cdot \frac{d\mathbf{r}}{dt} &= 2(t^2 \mathbf{i} + (t-1) \mathbf{j} + t^3 \mathbf{k}) \cdot (2t \mathbf{i} + \mathbf{j} + 3t^2 \mathbf{k}) \\ &= 2(2t^3 + (t-1) + 3t^5), \end{aligned}$$

or

$$\text{grad } u \cdot \frac{d\mathbf{r}}{dt} = \boxed{6t^5 + 4t^3 + 2t - 2},$$

which is the same as the expression for du/dt derived in Exercise 6.

0S.2.5 Derivatives “Following the Motion”

Solution to Exercise 1

The velocity at any point x in the centre of the stream at time t is $v(x, t)$. If the motion of the leaf (which is the same as that of the water immediately beneath it) is given by $x = x(t)$, its velocity at time t will be $v(x(t), t)$. But we can also find this velocity by differentiating the function $x(t)$. Thus

$$\frac{dx}{dt} = v(x(t), t).$$

Solution to Exercise 2

The temperature $T(t)$ at the leaf at time t will be equal to the temperature $T(x, t)$ at position x and time t , where $x = x(t)$. That is,

$$T(t) = T(x(t), t).$$

Solution to Exercise 3

(i) Since

$$T(x, t) = b \ln x + c \sin t,$$

and

$$x(t) = at + k,$$

we have

$$T(x(t), t) = b \ln(at + k) + c \sin t.$$

Thus, from the solution to Exercise 2,

$$T(t) = b \ln(at + k) + c \sin t.$$

(ii) Differentiating this expression,

$$\frac{dT}{dt} = \frac{ab}{at + k} + c \cos t.$$

(The first term of $T(t)$ can be differentiated using the chain rule for functions of one variable,

$$\frac{d}{dt}(b \ln x) = \frac{d}{dx}(b \ln x) \frac{dx}{dt},$$

where $x = at + k$.)

Solution to Exercise 4

Consider the data of Exercise 3. Computing the quantities to be included in the general expression for dT/dt we see that in Exercise 3 these were

$$\frac{\partial T}{\partial x} = \frac{b}{x} = \frac{b}{at + k},$$

$$\frac{\partial T}{\partial t} = c \cos t,$$

and

$$\frac{dx}{dt} = a.$$

Comparing with the solution to Exercise 3, we see that we can write

$$\frac{dT}{dt} = \boxed{\frac{dx}{dt} \frac{\partial T}{\partial x} + \frac{\partial T}{\partial t}}.$$

We could alternatively derive this result from the (boxed) chain rule in Exercise 4 of subsection 0.2.4 upon replacing u by T and y by t .

Solution to Exercise 5

The text of the exercise after the first sentence should read as follows:

“The symbol T on the left refers to the function $\boxed{T(t)}$ with domain \boxed{R} , describing the temperature at the leaf at time t . The symbol T on the right refers to the function $\boxed{T(x, t)}$, with domain $\boxed{R \times R}$, describing the temperature at position \boxed{x} at time t .”

Solution to Exercise 6

Replacing T by v in the solution to Exercise 4, we find

$$\frac{dv}{dt} = \boxed{\frac{dx}{dt} \frac{\partial v}{\partial x} + \frac{\partial v}{\partial t}}.$$

Solution to Exercise 7

The operator we seek is d/dt (as it appears on the left-hand sides of the solutions to Exercises 4 and 6). If we remove the particular dependent variable (v or T) involved in either of those expressions, we are left with

$$\frac{d}{dt} = \boxed{\frac{dx}{dt} \frac{\partial}{\partial x} + \frac{\partial}{\partial t}}.$$

It was shown in Exercise 1 that $dx/dt = v(x(t), t)$. Thus for the particular leaf, the operator can be written as

$$v(x(t), t) \frac{\partial}{\partial x} + \frac{\partial}{\partial t}.$$

We abandon the particular spot occupied by the leaf in favour of an arbitrary point of the stream by replacing the specific function $x(t)$ by the coordinate x . Writing v for $v(x, t)$, we are left with the operator equation

$$\frac{d}{dt} = \boxed{v \frac{\partial}{\partial x} + \frac{\partial}{\partial t}}.$$

0S.2.6 Change of Coordinates

Solution to Exercise 1

The expressions for x and y are

$$\begin{aligned} x &= \boxed{r \cos \theta}, \\ y &= \boxed{r \sin \theta}. \end{aligned}$$

Solution to Exercise 2

Since $T = f(x, y)$, we can use the formulae from Exercise 1 to get

$$T = \boxed{f(r \cos \theta, r \sin \theta)}.$$

Solution to Exercise 3

Along any curve for which $\theta = \text{constant}$ we have $T = f(x, y)$, where x and y are functions of r alone. Using the boxed formula

$$\boxed{\frac{du}{dt} = \frac{\partial u}{\partial x} \frac{dx}{dt} + \frac{\partial u}{\partial y} \frac{dy}{dt}}$$

(from Exercise 4 of subsection 0.2.4) with T for u and r for t , we have

$$\frac{dT}{dr} = \frac{\partial T}{\partial x} \frac{dx}{dr} + \frac{\partial T}{\partial y} \frac{dy}{dr} \text{ along } \theta = \text{constant}.$$

For functions of two variables r and θ however, the operator “ d/dr along $\theta = \text{constant}$ ” is, by definition, the same as $\partial/\partial r$. We can therefore write the above as

$$\frac{\partial T}{\partial r} = \frac{\partial T}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial T}{\partial y} \frac{\partial y}{\partial r}.$$

Since in this case

$$\frac{\partial x}{\partial r} = \cos \theta,$$

$$\frac{\partial y}{\partial r} = \sin \theta,$$

we have

$$\frac{\partial T}{\partial r} = \boxed{\cos \theta \frac{\partial T}{\partial x} + \sin \theta \frac{\partial T}{\partial y}}.$$

Similarly we can write

$$\frac{dT}{d\theta} = \frac{\partial T}{\partial x} \frac{dx}{d\theta} + \frac{\partial T}{\partial y} \frac{dy}{d\theta} \text{ along } r = \text{constant},$$

or

$$\frac{\partial T}{\partial \theta} = \frac{\partial T}{\partial x} \frac{\partial x}{\partial \theta} + \frac{\partial T}{\partial y} \frac{\partial y}{\partial \theta}.$$

Since

$$\frac{\partial x}{\partial \theta} = -r \sin \theta,$$

$$\frac{\partial y}{\partial \theta} = r \cos \theta,$$

we arrive at

$$\frac{\partial T}{\partial \theta} = \boxed{-r \sin \theta \frac{\partial T}{\partial x} + r \cos \theta \frac{\partial T}{\partial y}}.$$

Solution to Exercise 4

Since $T = f(x, y)$, and $x = \alpha p + \beta q$, $y = \gamma p + \delta q$,

$$T = \boxed{f(\alpha p + \beta q, \gamma p + \delta q)}.$$

We obtain the expressions for $\partial T/\partial p$ and $\partial T/\partial q$ as before by using the boxed formula

$$\boxed{\frac{du}{dt} = \frac{\partial u}{\partial x} \frac{dx}{dt} + \frac{\partial u}{\partial y} \frac{dy}{dt}}$$

(from Exercise 4 of subsection 0.2.4) with the total derivatives replaced by partial derivatives, T for u , and first p and then q in place of t :

$$\frac{\partial T}{\partial p} = \frac{\partial T}{\partial x} \frac{\partial x}{\partial p} + \frac{\partial T}{\partial y} \frac{\partial y}{\partial p},$$

$$\frac{\partial T}{\partial q} = \frac{\partial T}{\partial x} \frac{\partial x}{\partial q} + \frac{\partial T}{\partial y} \frac{\partial y}{\partial q}.$$

Since

$$\frac{\partial x}{\partial p} = \alpha, \frac{\partial y}{\partial p} = \gamma, \frac{\partial x}{\partial q} = \beta, \frac{\partial y}{\partial q} = \delta,$$

we have

$$\frac{\partial T}{\partial p} = \boxed{\alpha \frac{\partial T}{\partial x} + \gamma \frac{\partial T}{\partial y}},$$

$$\frac{\partial T}{\partial q} = \boxed{\beta \frac{\partial T}{\partial x} + \delta \frac{\partial T}{\partial y}}.$$

Solution to Exercise 5

We proceed as before. Since $T = f(x, y)$ and $x = x(\xi, \eta)$, $y = y(\xi, \eta)$,

$$T = \boxed{f(x(\xi, \eta), y(\xi, \eta))}.$$

The chain rule is used as in Exercises 3 and 4 to give

$$\frac{\partial T}{\partial \xi} = \boxed{\frac{\partial T}{\partial x} \frac{\partial x}{\partial \xi} + \frac{\partial T}{\partial y} \frac{\partial y}{\partial \xi}},$$

$$\frac{\partial T}{\partial \eta} = \boxed{\frac{\partial T}{\partial x} \frac{\partial x}{\partial \eta} + \frac{\partial T}{\partial y} \frac{\partial y}{\partial \eta}}.$$

Solution to Exercise 6

The symbols used here are differently arranged from those in the general case of Exercise 5. We are given $\xi = \xi(x, t)$, $\eta = \eta(x, t)$ (for particular functions ξ and η), and the corresponding chain rule is

$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial \xi} \frac{\partial \xi}{\partial x} + \frac{\partial u}{\partial \eta} \frac{\partial \eta}{\partial x},$$

$$\frac{\partial u}{\partial t} = \frac{\partial u}{\partial \xi} \frac{\partial \xi}{\partial t} + \frac{\partial u}{\partial \eta} \frac{\partial \eta}{\partial t}.$$

Since

$$\xi(x, t) = x + ct,$$

$$\eta(x, t) = x - ct,$$

we have

$$\frac{\partial \xi}{\partial x} = \frac{\partial \eta}{\partial x} = 1,$$

$$\frac{\partial \xi}{\partial t} = -\frac{\partial \eta}{\partial t} = c,$$

so that

$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial \xi} + \frac{\partial u}{\partial \eta},$$

$$\frac{\partial u}{\partial t} = c \left(\frac{\partial u}{\partial \xi} - \frac{\partial u}{\partial \eta} \right).$$

Solution to Exercise 7

We have $T(x, z) = T(x, y) = f(x, y)$ where $y = x + z$. Using the general form of the chain rule appearing in the last exercise, with u, t, ξ, η being replaced by $T(x, z) = f(x, y)$, z, x and y respectively,

$$\frac{\partial T}{\partial x}(x, z) = \frac{\partial f}{\partial x} + \frac{\partial y}{\partial x} \frac{\partial f}{\partial y},$$

$$\frac{\partial T}{\partial z}(x, z) = \frac{\partial y}{\partial z} \frac{\partial f}{\partial y}$$

(since $\partial x / \partial x = 1$ and $\partial x / \partial z = 0$). Thus

$$\frac{\partial T}{\partial x}(x, z) = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y},$$

$$\frac{\partial T}{\partial z}(x, z) = \frac{\partial f}{\partial y},$$

or

$$\frac{\partial T}{\partial x}(x, z) = \frac{\partial T}{\partial x}(x, y) + \frac{\partial T}{\partial y}(x, y),$$

$$\frac{\partial T}{\partial z}(x, z) = \frac{\partial T}{\partial y}(x, y).$$

Solution to Exercise 8

By putting u for T in the solution to Exercise 7 we see that in terms of the variables x, z the equation takes the form

$$\frac{\partial u}{\partial x}(x, z) = 0.$$

Solution to Exercise 9

We see from Exercise 2 of subsection 0.2.2 that the general solution of

$$\frac{\partial u}{\partial x}(x, z) = 0$$

is

$$u(x, z) = f(z),$$

where f is an arbitrary function of one variable. Since $y = x + z$, we have $z = y - x$, so that the general solution of

$$\frac{\partial u}{\partial x}(x, y) + \frac{\partial u}{\partial y}(x, y) = 0$$

is

$$u(x, y) = f(y - x),$$

which agrees with the result suggested by Exercise 2 of subsection 0.2.3.

Solution to Exercise 10

We are asked to restate the general form of the chain rule given in the solution to Exercise 6 (with t being replaced by y) in two different ways.

(i) As operator equations these are

$$\begin{aligned} \frac{\partial}{\partial x} &= \frac{\partial \xi}{\partial x} \frac{\partial}{\partial \xi} + \frac{\partial \eta}{\partial x} \frac{\partial}{\partial \eta}, \\ \frac{\partial}{\partial y} &= \frac{\partial \xi}{\partial y} \frac{\partial}{\partial \xi} + \frac{\partial \eta}{\partial y} \frac{\partial}{\partial \eta}. \end{aligned}$$

(ii) As a matrix equation, they can be expressed by

$$\begin{bmatrix} \frac{\partial u}{\partial x} \\ \frac{\partial u}{\partial y} \end{bmatrix} = \begin{bmatrix} \frac{\partial \xi}{\partial x} & \frac{\partial \eta}{\partial x} \\ \frac{\partial \xi}{\partial y} & \frac{\partial \eta}{\partial y} \end{bmatrix} \begin{bmatrix} \frac{\partial u}{\partial \xi} \\ \frac{\partial u}{\partial \eta} \end{bmatrix}.$$

Solution to Exercise 11

There are two methods of solution. The first involves performing the matrix multiplication explicitly and using the chain rule:

$$\begin{aligned}
 AB &= \begin{bmatrix} \frac{\partial \xi}{\partial x} & \frac{\partial \eta}{\partial x} & \frac{\partial x}{\partial \xi} & \frac{\partial y}{\partial \xi} \\ \frac{\partial \xi}{\partial y} & \frac{\partial \eta}{\partial y} & \frac{\partial x}{\partial \eta} & \frac{\partial y}{\partial \eta} \end{bmatrix} \\
 &= \begin{bmatrix} \left(\frac{\partial \xi}{\partial x} \frac{\partial x}{\partial \xi} + \frac{\partial \eta}{\partial x} \frac{\partial x}{\partial \eta} \right) & \left(\frac{\partial \xi}{\partial x} \frac{\partial y}{\partial \xi} + \frac{\partial \eta}{\partial x} \frac{\partial y}{\partial \eta} \right) \\ \left(\frac{\partial \xi}{\partial y} \frac{\partial x}{\partial \xi} + \frac{\partial \eta}{\partial y} \frac{\partial x}{\partial \eta} \right) & \left(\frac{\partial \xi}{\partial y} \frac{\partial y}{\partial \xi} + \frac{\partial \eta}{\partial y} \frac{\partial y}{\partial \eta} \right) \end{bmatrix},
 \end{aligned}$$

Using the solutions to Exercise 10(i) we see that this is equal to

$$\begin{bmatrix} \frac{\partial x}{\partial x} & \frac{\partial y}{\partial x} \\ \frac{\partial x}{\partial y} & \frac{\partial y}{\partial y} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix},$$

so that

$$AB = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

The second method is to note that corresponding to the matrix equation of Exercise 10(ii) giving $\partial u/\partial x$, $\partial u/\partial y$ in terms of $\partial u/\partial \xi$ and $\partial u/\partial \eta$, there will be a similar equation giving $\partial u/\partial \xi$, $\partial u/\partial \eta$ in terms of $\partial u/\partial x$ and $\partial u/\partial y$. The chain rule says that this equation is

$$\begin{bmatrix} \frac{\partial u}{\partial \xi} \\ \frac{\partial u}{\partial \eta} \end{bmatrix} = \begin{bmatrix} \frac{\partial x}{\partial \xi} & \frac{\partial y}{\partial \xi} \\ \frac{\partial x}{\partial \eta} & \frac{\partial y}{\partial \eta} \end{bmatrix} \begin{bmatrix} \frac{\partial u}{\partial x} \\ \frac{\partial u}{\partial y} \end{bmatrix}$$

and the 2×2 matrix here is just B . By combining this equation with that of Exercise 10(ii) we see immediately that A and B must be inverse to each other, so that AB is the identity matrix. This result is the analogue in two dimensions of the one-dimensional formula

$$\frac{dy}{dx} \frac{dx}{dy} = 1;$$

but note that, for example,

$$\frac{\partial x}{\partial \xi} \frac{\partial \xi}{\partial x} \neq 1$$

in general, since the $(1, 1)$ elements of two matrices inverse to each other are not normally reciprocals.

0S.3 SECOND-ORDER PARTIAL DIFFERENTIATION

0S.3.1 Second and Higher Order Partial Derivatives

Solution to Exercise 1

If $u(x, t) = \sin xt + \sin x \sin t + t$ then

$$\begin{aligned}\frac{\partial u}{\partial x} &= t \cos xt + \cos x \sin t \\ \frac{\partial u}{\partial t} &= x \cos xt + \sin x \cos t + 1 \\ \frac{\partial^2 u}{\partial x \partial t} &= \cos xt - xt \sin xt + \cos x \cos t \\ \frac{\partial^2 u}{\partial t \partial x} &= \cos xt - xt \sin xt + \cos x \cos t \\ \frac{\partial^2 u}{\partial t^2} &= -x^2 \sin xt - \sin x \sin t\end{aligned}$$

Solution to Exercise 2

Yes,

provided that the two mixed derivatives exist and are continuous. All the functions to be encountered in the course will satisfy these conditions.

Solution to Exercise 3

Since $\partial^2/\partial x^2$ and $\partial^2/\partial y^2$ commute, we have

$$\begin{aligned}\frac{\partial^{2n} f}{\partial x^{2n}} &= \left(\frac{\partial^2}{\partial x^2} \right)^n f \\ &= \left(\frac{\partial^2}{\partial x^2} \right)^{n-1} \left(\frac{\partial^2}{\partial x^2} \right) f \\ &= \left(\frac{\partial^2}{\partial x^2} \right)^{n-1} \left(\frac{\partial^2}{\partial y^2} \right) f \\ &= \frac{\partial^2}{\partial y^2} \left(\frac{\partial^2}{\partial x^2} \right)^{n-1} f \\ &= \frac{\partial^2}{\partial y^2} \left(\frac{\partial^2}{\partial x^2} \right)^{n-2} \left(\frac{\partial^2}{\partial y^2} \right) f \\ &= \left(\frac{\partial^2}{\partial y^2} \right)^2 \left(\frac{\partial^2}{\partial x^2} \right)^{n-2} f,\end{aligned}$$

and so on until all the $\partial^2/\partial x^2$ have been changed into $\partial^2/\partial y^2$. Hence the result is

$$\left(\frac{\partial^2}{\partial x^2} \right)^n f = \left(\frac{\partial^2}{\partial y^2} \right)^n f,$$

or

$$\frac{\partial^{2n} f}{\partial x^{2n}} = \frac{\partial^{2n} f}{\partial y^{2n}}.$$

Solution to Exercise 4

This time we have

$$\left(\frac{\partial^2}{\partial x^2}\right)f = \left(-\frac{\partial^2}{\partial y^2}\right)f,$$

and so, the same procedure as we used in the preceding exercise gives

$$\left(\frac{\partial^2}{\partial x^2}\right)^n f = \left(-\frac{\partial^2}{\partial y^2}\right)^n f,$$

which can be written as

$$\frac{\partial^{2n} f}{\partial x^{2n}} = \boxed{(-1)^n \frac{\partial^{2n} f}{\partial y^{2n}}}.$$

Solution to Exercise 5

Proceeding as before we find that

$$\frac{\partial^n f}{\partial y^n} = \boxed{\frac{\partial^{2n} f}{\partial x^{2n}}}.$$

Hence

$$\frac{\partial^2 f}{\partial y^2} - 3 \frac{\partial^3 f}{\partial y \partial x^2} + 2 \frac{\partial^4 f}{\partial x^4} = \frac{\partial^2 f}{\partial y^2} - 3 \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial y} \right) + 2 \frac{\partial^2 f}{\partial y^2},$$

giving

$$\frac{\partial^2 f}{\partial y^2} - 3 \frac{\partial^3 f}{\partial y \partial x^2} + 2 \frac{\partial^4 f}{\partial x^4} = \boxed{0}.$$

0S.3.2 Solving Simple Second-Order Partial Differential Equations**Solution to Exercise 1**

Putting $\partial u / \partial y = p$, the equation becomes

$$\boxed{\frac{\partial p}{\partial x} = 0},$$

with solution

$$p(x, y) = \boxed{f(y)}$$

for an arbitrary function f . Since

$$\frac{\partial u}{\partial y}(x, y) = \boxed{f(y)}$$

has solution

$$u = \int f(y) + g(x),$$

the general solution of the original equation is

$$u(x, y) = \int f(y) + g(x)$$

where f and g are arbitrary functions of one variable. Since the equation we started from is symmetric in x and y we could equally well have solved the problem by putting $\partial u / \partial x = p$ and following the same method thereafter. We would then have arrived at the general solution

$$u(x, y) = h(y) + \int g_1(x),$$

where h, g_1 are arbitrary. It follows from this that we can express the general solution most simply as

$$u(x, y) = g(x) + h(y).$$

Since h is a primitive of f and g is a primitive of g_1 , both h and g must be differentiable, but are otherwise arbitrary.

Solution to Exercise 2

Putting $\partial u / \partial x = p$, the equation becomes

$$\frac{\partial p}{\partial x} = 0,$$

with solution

$$p(x, y) = f(y),$$

for arbitrary f . Since the equation

$$\frac{\partial u}{\partial x} = f(y)$$

has solution

$$u = xf(y) + g(y),$$

we have for the general solution of the original equation

$$u(x, y) = g(y) + xf(y),$$

where f and g are arbitrary functions of one variable.

Solution to Exercise 3

Putting $\partial u / \partial y = p$, the equation becomes

$$\frac{\partial p}{\partial y} - \frac{1}{y}p = 0.$$

An integrating factor is

$$\begin{aligned} \exp \int \left(-\frac{1}{y} \right) dy &= \exp(-\ln y) \\ &= \frac{1}{y} \end{aligned}$$

(compare with Exercise 5 of subsection 0.2.2), and multiplying through by this factor brings the equation to the form

$$\frac{\partial}{\partial y} \left(\frac{p(x, y)}{y} \right) = 0,$$

which has solution

$$p(x, y) = \boxed{yf(x)},$$

where f is arbitrary. Since $p = \partial u / \partial y$, an integration with respect to y gives

$$u(x, y) = \boxed{\frac{1}{2}y^2f(x) + g(x)}.$$

The factor $\frac{1}{2}$ may without loss of generality be absorbed into the arbitrary function f .

Solution to Exercise 4

In terms of p , where $p = \partial u / \partial y$, the equation is

$$x \frac{\partial p}{\partial x} = 2p,$$

or in normal form,

$$\frac{\partial p}{\partial x} - \frac{2}{x}p = 0.$$

An integrating factor is

$$\exp \int \left(-\frac{2}{x} \right) dx = \frac{1}{x^2},$$

bringing the equation to the form

$$\frac{\partial}{\partial x} \left(\frac{p(x, y)}{x^2} \right) = 0.$$

Integrating with respect to x and solving for p gives

$$p(x, y) = x^2 f(y)$$

with f arbitrary. Since $p = \partial u / \partial y$ we can now integrate with respect to y to obtain

$$u(x, y) = \boxed{x^2 \int f(y) + g(x)}.$$

0S.3.3 Transforming Second-Order Derivatives

Solution to Exercise 1

From Exercise 6 of subsection 0.2.6,

$$\frac{\partial}{\partial x} = \frac{\partial}{\partial \xi} + \frac{\partial}{\partial \eta},$$

$$\frac{\partial}{\partial t} = c \left(\frac{\partial}{\partial \xi} - \frac{\partial}{\partial \eta} \right).$$

Since $\partial^2/\partial x^2 = (\partial/\partial x)^2$, $\partial^2/\partial x \partial t = (\partial/\partial x)(\partial/\partial t)$ etc., we then have

$$\frac{\partial^2}{\partial x^2} = \frac{\partial^2}{\partial \xi^2} + 2 \frac{\partial^2}{\partial \xi \partial \eta} + \frac{\partial^2}{\partial \eta^2},$$

$$\frac{\partial^2}{\partial x \partial t} = c \left(\frac{\partial^2}{\partial \xi^2} - \frac{\partial^2}{\partial \eta^2} \right),$$

$$\frac{\partial^2}{\partial t^2} = c^2 \left(\frac{\partial^2}{\partial \xi^2} - 2 \frac{\partial^2}{\partial \xi \partial \eta} + \frac{\partial^2}{\partial \eta^2} \right).$$

Solution to Exercise 2

We see from Exercise 1 that

$$\frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 u}{\partial \xi^2} + 2 \frac{\partial^2 u}{\partial \xi \partial \eta} + \frac{\partial^2 u}{\partial \eta^2},$$

$$\frac{\partial^2 u}{\partial t^2} = c^2 \left(\frac{\partial^2 u}{\partial \xi^2} - 2 \frac{\partial^2 u}{\partial \xi \partial \eta} + \frac{\partial^2 u}{\partial \eta^2} \right).$$

Subtracting the second equation from c^2 times the first, we have

$$c^2 \frac{\partial^2 u}{\partial x^2} - \frac{\partial^2 u}{\partial t^2} = 4c^2 \frac{\partial^2 u}{\partial \xi \partial \eta},$$

so that

$$\frac{\partial^2 u}{\partial \xi \partial \eta} = 0.$$

By reference to the solution of Exercise 1 in Subsection 0.3.2 we see that the general solution of this equation is

$$u(\xi, \eta) = f(\xi) + g(\eta),$$

for arbitrary differentiable functions f and g , so that the solution of the wave equation is

$$u(x, t) = f(x + ct) + g(x - ct).$$

Solution to Exercise 3

Using the chain rule,

$$\frac{\partial}{\partial x} = \frac{\partial \xi}{\partial x} \frac{\partial}{\partial \xi} + \frac{\partial \eta}{\partial x} \frac{\partial}{\partial \eta},$$

$$\frac{\partial}{\partial t} = \frac{\partial \xi}{\partial t} \frac{\partial}{\partial \xi} + \frac{\partial \eta}{\partial t} \frac{\partial}{\partial \eta}.$$

Since the given expressions for ξ, η mean that

$$\frac{\partial \xi}{\partial x} = \alpha, \frac{\partial \xi}{\partial t} = \beta, \frac{\partial \eta}{\partial x} = \gamma, \frac{\partial \eta}{\partial t} = \delta,$$

we have

$$\frac{\partial}{\partial x} = \alpha \frac{\partial}{\partial \xi} + \gamma \frac{\partial}{\partial \eta},$$

$$\frac{\partial}{\partial t} = \beta \frac{\partial}{\partial \xi} + \delta \frac{\partial}{\partial \eta}.$$

The second-order derivatives are, therefore,

$$\frac{\partial^2}{\partial x^2} = \alpha^2 \frac{\partial^2}{\partial \xi^2} + 2\alpha\gamma \frac{\partial^2}{\partial \xi \partial \eta} + \gamma^2 \frac{\partial^2}{\partial \eta^2},$$

$$\frac{\partial^2}{\partial x \partial t} = \alpha\beta \frac{\partial^2}{\partial \xi^2} + (\alpha\delta + \beta\gamma) \frac{\partial^2}{\partial \xi \partial \eta} + \gamma\delta \frac{\partial^2}{\partial \eta^2},$$

$$\frac{\partial^2}{\partial t^2} = \beta^2 \frac{\partial^2}{\partial \xi^2} + 2\beta\delta \frac{\partial^2}{\partial \xi \partial \eta} + \delta^2 \frac{\partial^2}{\partial \eta^2},$$

so that

$$L = \left(A\beta^2 + B\alpha\beta + C\alpha^2 \right) \frac{\partial^2}{\partial \xi^2} + (2A\beta\delta + B(\alpha\delta + \beta\gamma) + 2C\alpha\gamma) \frac{\partial^2}{\partial \xi \partial \eta} + (A\delta^2 + B\gamma\delta + C\gamma^2) \frac{\partial^2}{\partial \eta^2}$$

Solution to Exercise 4

By either of the ways suggested we get

$$\frac{\partial}{\partial r} = \cos \theta \frac{\partial}{\partial x} + \sin \theta \frac{\partial}{\partial y},$$

$$\frac{\partial}{\partial \theta} = -r \sin \theta \frac{\partial}{\partial x} + r \cos \theta \frac{\partial}{\partial y}.$$

Solution to Exercise 5

We have

$$\begin{bmatrix} \frac{\partial}{\partial r} \\ \frac{\partial}{\partial \theta} \end{bmatrix} = \begin{bmatrix} \cos \theta & \sin \theta \\ -r \sin \theta & r \cos \theta \end{bmatrix} \begin{bmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \end{bmatrix}.$$

The matrix of coefficients can be inverted in the usual way:

$$\begin{bmatrix} \cos \theta & \sin \theta & 1 & 0 \\ -r \sin \theta & r \cos \theta & 0 & 1 \end{bmatrix}$$

Stage 1: Multiply first row by $\sec \theta$ and then add $r \sin \theta$ times the first row to the second row. This gives

$$\begin{bmatrix} 1 & \tan \theta & \sec \theta & 0 \\ 0 & r \sec \theta & r \tan \theta & 1 \end{bmatrix}.$$

Stage 2: Multiply second row by $(1/r) \cos \theta$ and then subtract $\tan \theta$ times the second row from the first row. This gives

$$\begin{bmatrix} 1 & 0 & \cos \theta & -\frac{1}{r} \sin \theta \\ 0 & 1 & \sin \theta & \frac{1}{r} \cos \theta \end{bmatrix}.$$

Reading off the inverse of the matrix we started with,

$$\begin{bmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \end{bmatrix} = \begin{bmatrix} \cos \theta & -\frac{1}{r} \sin \theta \\ \sin \theta & \frac{1}{r} \cos \theta \end{bmatrix} \begin{bmatrix} \frac{\partial}{\partial r} \\ \frac{\partial}{\partial \theta} \end{bmatrix},$$

or in other words,

$$\begin{aligned} \frac{\partial}{\partial x} &= \cos \theta \frac{\partial}{\partial r} - \frac{1}{r} \sin \theta \frac{\partial}{\partial \theta}, \\ \frac{\partial}{\partial y} &= \sin \theta \frac{\partial}{\partial r} + \frac{1}{r} \cos \theta \frac{\partial}{\partial \theta}. \end{aligned}$$

Solution to Exercise 6

$$\begin{aligned} \frac{\partial^2}{\partial x^2} &= \left(\cos \theta \frac{\partial}{\partial r} - \frac{1}{r} \sin \theta \frac{\partial}{\partial \theta} \right) \left(\cos \theta \frac{\partial}{\partial r} - \frac{1}{r} \sin \theta \frac{\partial}{\partial \theta} \right) \\ &= \cos^2 \theta \frac{\partial^2}{\partial r^2} - \cos \theta \frac{\partial}{\partial r} \left(\frac{1}{r} \sin \theta \frac{\partial}{\partial \theta} \right) - \frac{1}{r} \sin \theta \frac{\partial}{\partial \theta} \left(\cos \theta \frac{\partial}{\partial r} \right) \\ &\quad + \frac{1}{r} \sin \theta \frac{\partial}{\partial \theta} \left(\frac{1}{r} \sin \theta \frac{\partial}{\partial \theta} \right) \\ &= \cos^2 \theta \frac{\partial^2}{\partial r^2} + \frac{\sin \theta \cos \theta}{r^2} \frac{\partial}{\partial \theta} - \frac{\sin \theta \cos \theta}{r} \frac{\partial^2}{\partial r \partial \theta} + \frac{\sin^2 \theta}{r} \frac{\partial}{\partial r} \\ &\quad - \frac{\sin \theta \cos \theta}{r} \frac{\partial^2}{\partial r \partial \theta} + \frac{\sin \theta \cos \theta}{r^2} \frac{\partial}{\partial \theta} + \frac{\sin^2 \theta}{r^2} \frac{\partial^2}{\partial \theta^2}. \end{aligned}$$

Therefore

$$\frac{\partial^2}{\partial x^2} = \cos^2 \theta \frac{\partial^2}{\partial r^2} - \frac{2 \sin \theta \cos \theta}{r} \frac{\partial^2}{\partial r \partial \theta} + \frac{\sin^2 \theta}{r^2} \frac{\partial^2}{\partial \theta^2} + \frac{\sin^2 \theta}{r} \frac{\partial}{\partial r} + \frac{2 \sin \theta \cos \theta}{r^2} \frac{\partial}{\partial \theta}.$$

Similarly,

$$\begin{aligned} \frac{\partial^2}{\partial y^2} &= \left(\sin \theta \frac{\partial}{\partial r} + \frac{1}{r} \cos \theta \frac{\partial}{\partial \theta} \right) \left(\sin \theta \frac{\partial}{\partial r} + \frac{1}{r} \cos \theta \frac{\partial}{\partial \theta} \right) \\ &= \sin^2 \theta \frac{\partial^2}{\partial r^2} + \sin \theta \frac{\partial}{\partial r} \left(\frac{1}{r} \cos \theta \frac{\partial}{\partial \theta} \right) + \frac{1}{r} \cos \theta \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial r} \right) \\ &\quad + \frac{1}{r} \cos \theta \frac{\partial}{\partial \theta} \left(\frac{1}{r} \cos \theta \frac{\partial}{\partial \theta} \right) \end{aligned}$$

$$\begin{aligned}
&= \sin^2 \theta \frac{\partial^2}{\partial r^2} - \frac{\sin \theta \cos \theta}{r^2} \frac{\partial}{\partial \theta} + \frac{\sin \theta \cos \theta}{r} \frac{\partial^2}{\partial r \partial \theta} + \frac{\cos^2 \theta}{r} \frac{\partial}{\partial r} \\
&\quad + \frac{\sin \theta \cos \theta}{r} \frac{\partial^2}{\partial r \partial \theta} - \frac{\sin \theta \cos \theta}{r^2} \frac{\partial}{\partial \theta} + \frac{\cos^2 \theta}{r^2} \frac{\partial^2}{\partial \theta^2}.
\end{aligned}$$

Therefore,

$$\frac{\partial^2}{\partial y^2} = \left[\sin^2 \theta \frac{\partial^2}{\partial r^2} + \frac{2 \sin \theta \cos \theta}{r} \frac{\partial^2}{\partial r \partial \theta} + \frac{\cos^2 \theta}{r^2} \frac{\partial^2}{\partial \theta^2} + \frac{\cos^2 \theta}{r} \frac{\partial}{\partial r} - \frac{2 \sin \theta \cos \theta}{r^2} \frac{\partial}{\partial \theta} \right].$$

It follows that

$$\nabla^2 = \left[\frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} + \frac{1}{r^2} \frac{\partial^2}{\partial \theta^2} \right].$$

Solution to Exercise 7

(i) Since

$$\frac{\partial \xi}{\partial x} = 2x, \frac{\partial \eta}{\partial x} = 0, \frac{\partial \xi}{\partial t} = 2t, \frac{\partial \eta}{\partial t} = 1,$$

we have by the chain rule,

$$\frac{\partial}{\partial x} = \left[2x \frac{\partial}{\partial \xi} \right],$$

$$\frac{\partial}{\partial t} = \left[2t \frac{\partial}{\partial \xi} + \frac{\partial}{\partial \eta} \right].$$

(ii) From the first answer in part (i), we have

$$\frac{\partial}{\partial \eta} = \left[\frac{1}{2x} \frac{\partial}{\partial x} \right].$$

Substituting this into the second answer in part (i) gives

$$\frac{\partial}{\partial \eta} = \left[\frac{\partial}{\partial t} - \frac{t}{x} \frac{\partial}{\partial x} \right].$$

(iii)

$$\begin{aligned}
\frac{\partial^2}{\partial \eta^2} &= \left(\frac{\partial}{\partial t} - \frac{t}{x} \frac{\partial}{\partial x} \right) \left(\frac{\partial}{\partial t} - \frac{t}{x} \frac{\partial}{\partial x} \right) \\
&= \frac{\partial^2}{\partial t^2} - \frac{\partial}{\partial t} \left(\frac{t}{x} \frac{\partial}{\partial x} \right) - \frac{t}{x} \frac{\partial^2}{\partial x \partial t} + \frac{t}{x} \frac{\partial}{\partial x} \left(\frac{t}{x} \frac{\partial}{\partial x} \right) \\
&= \frac{\partial^2}{\partial t^2} - \frac{1}{x} \frac{\partial}{\partial x} - \frac{t}{x} \frac{\partial^2}{\partial x \partial t} - \frac{t}{x} \frac{\partial^2}{\partial x \partial t} - \frac{t^2}{x^3} \frac{\partial}{\partial x} + \frac{t^2}{x^2} \frac{\partial^2}{\partial x^2},
\end{aligned}$$

so that we have

$$\frac{\partial^2}{\partial \eta^2} = \frac{1}{x^2} L - \left[\frac{1}{x} \left(1 + \frac{t^2}{x^2} \right) \frac{\partial}{\partial x} \right].$$

(iv) The last equation in part (iii) can be written as

$$L = x^2 \frac{\partial^2}{\partial \eta^2} + x \left(1 + \frac{t^2}{x^2} \right) \frac{\partial}{\partial x},$$

and using the first answer in part (i) gives

$$L = \boxed{x^2 \frac{\partial^2}{\partial \eta^2} + 2(x^2 + t^2) \frac{\partial}{\partial \xi}}.$$

Solution to Exercise 8

Using the solution to Exercise 7(i), we see that the right-hand side of the given equation can be expressed as

$$\frac{x^2}{t} \left(2t \frac{\partial u}{\partial \xi} + \frac{\partial u}{\partial \eta} \right) + \frac{t^2}{x} \left(2x \frac{\partial u}{\partial \xi} \right) = 2(x^2 + t^2) \frac{\partial u}{\partial \xi} + \frac{x^2}{t} \frac{\partial u}{\partial \eta},$$

so that upon using the expression for L derived in Exercise 7(iv), dividing through by x^2 and replacing t by η , the transformed equation is seen to be

$$\boxed{\frac{\partial^2 u}{\partial \eta^2} = \frac{1}{\eta} \frac{\partial u}{\partial \eta}}.$$

By reference to Exercise 3 of subsection 0.3.2 we see that the general solution of the transformed equation is

$$u(\xi, \eta) = \boxed{f(\xi) + \eta^2 g(\xi)},$$

where f and g are arbitrary functions of one variable. Thus the general solution of the original equation is

$$u(x, t) = \boxed{f(x^2 + t^2) + t^2 g(x^2 + t^2)}.$$

Solution to Exercise 9

In Exercise 7 of subsection 0.1.1 we saw that

$$\frac{d}{dx}(\arctan x) = \frac{1}{1 + x^2},$$

and using this result in conjunction with the composite function rule we get

$$\frac{d}{dx} \left(\frac{1}{2} \arctan \frac{1}{2} x \right) = \frac{1}{4 + x^2},$$

so that we have

$$\frac{\partial \xi}{\partial x} = \frac{1}{1 + x^2}, \quad \frac{\partial \xi}{\partial t} = -1,$$

$$\frac{\partial \eta}{\partial x} = \frac{1}{4 + x^2}, \quad \frac{\partial \eta}{\partial t} = -1.$$

The chain rule then gives

$$\frac{\partial}{\partial x} = \frac{1}{1 + x^2} \frac{\partial}{\partial \xi} + \frac{1}{4 + x^2} \frac{\partial}{\partial \eta},$$

$$\frac{\partial}{\partial t} = - \left(\frac{\partial}{\partial \xi} + \frac{\partial}{\partial \eta} \right).$$

Solving these equations for $\partial/\partial \xi$ and $\partial/\partial \eta$ we find

$$\frac{\partial}{\partial \xi} = \frac{1}{3}(1 + x^2)(4 + x^2) \left(\frac{\partial}{\partial x} + \frac{1}{4 + x^2} \frac{\partial}{\partial t} \right),$$

$$\frac{\partial}{\partial \eta} = -\frac{1}{3}(1 + x^2)(4 + x^2) \left(\frac{\partial}{\partial x} + \frac{1}{1 + x^2} \frac{\partial}{\partial t} \right).$$

It follows that

$$\begin{aligned}
 \frac{\partial^2}{\partial \xi \partial \eta} &= -\frac{1}{9}(1+x^2)(4+x^2) \left(\frac{\partial}{\partial x} + \frac{1}{4+x^2} \frac{\partial}{\partial t} \right) \\
 &\quad \times \left\{ (1+x^2)(4+x^2) \left(\frac{\partial}{\partial x} + \frac{1}{1+x^2} \frac{\partial}{\partial t} \right) \right\} \\
 &= -\frac{1}{9}(1+x^2)(4+x^2) \left\{ \frac{\partial^2}{\partial t^2} + (5+2x^2) \frac{\partial^2}{\partial x \partial t} + (1+x^2)(4+x^2) \frac{\partial^2}{\partial x^2} \right. \\
 &\quad \left. + 2x(5+2x^2) \frac{\partial}{\partial x} + 2x \frac{\partial}{\partial t} \right\} \\
 &= -\frac{1}{9}(1+x^2)(4+x^2) \left\{ L + 2x \left[(5+2x^2) \frac{\partial}{\partial x} + \frac{\partial}{\partial t} \right] \right\}.
 \end{aligned}$$

Using the expressions above for $\partial/\partial x$, $\partial/\partial t$ in terms of $\partial/\partial \xi$, $\partial/\partial \eta$,

$$(5+2x^2) \frac{\partial}{\partial x} + \frac{\partial}{\partial t} = \frac{4+x^2}{1+x^2} \frac{\partial}{\partial \xi} + \frac{1+x^2}{4+x^2} \frac{\partial}{\partial \eta},$$

so that the expression for L is

$$L = \boxed{\frac{-9}{(1+x^2)(4+x^2)} \left[\frac{\partial^2}{\partial \xi \partial \eta} + \frac{2x}{9}(4+x^2)^2 \frac{\partial}{\partial \xi} + \frac{2x}{9}(1+x^2)^2 \frac{\partial}{\partial \eta} \right]},$$

which is equivalent to the formula for $L[u]$ given at the top of page 46 in W .

0S.4 VECTOR ALGEBRA AND LINE INTEGRALS

0S.4.1 Vector Algebra

Solution to Exercise 1

- (i) If we complete the parallelogram which has (arrows belonging to) \mathbf{a} and \mathbf{b} as adjacent sides, then the diagonal from the point O where the sides \mathbf{a} and \mathbf{b} meet is $\mathbf{a} + \mathbf{b}$. Since the diagonals of a parallelogram bisect each other, $\frac{1}{2}(\mathbf{a} + \mathbf{b})$ will be the position vector of a point halfway between the two vertices of the parallelogram with position vectors \mathbf{a} and \mathbf{b} (taking O as the origin). The answer is therefore

$$\frac{1}{2}(\mathbf{a} + \mathbf{b}) .$$

- (ii) The answer here depends on how the arrows are arranged. If \mathbf{b} points from the end of the arrow \mathbf{a} (or vice versa) then the third side of the triangle is

$$\mathbf{a} + \mathbf{b} ,$$

or $-\mathbf{a} - \mathbf{b}$, depending upon how the vector is directed along the side. If \mathbf{a} and \mathbf{b} have corresponding ends coincident then the third side of the triangle will be $\mathbf{a} - \mathbf{b}$ or $\mathbf{b} - \mathbf{a}$, depending on what direction is associated with it.

- (iii) Since by the definition of the scalar product,

$$\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}| |\mathbf{b}| \cos \theta ,$$

where θ is the angle between \mathbf{a} and \mathbf{b} , the required cosine is equal to

$$\frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}| |\mathbf{b}|} .$$

- (iv) The projection of \mathbf{a} along the direction of \mathbf{b} is

$$|\mathbf{a}| \cos \theta ,$$

which, we see from part (iii), is equal to

$$\frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{b}|} .$$

- (v) Since $\mathbf{a} \times \mathbf{b}$ is perpendicular to both \mathbf{a} and \mathbf{b} the required unit vector is

$$\frac{\mathbf{a} \times \mathbf{b}}{|\mathbf{a} \times \mathbf{b}|} .$$

- (vi) The parallelogram's area is $|\mathbf{a}| |\mathbf{b}| \sin \theta$, but we have $|\mathbf{a} \times \mathbf{b}| = |\mathbf{a}| |\mathbf{b}| \sin \theta$ by definition. The area is therefore

$$|\mathbf{a} \times \mathbf{b}| .$$

- (vii) \mathbf{a} and \mathbf{b} are perpendicular if $\theta = \pi/2$, or $\cos \theta = 0$, or

$$\mathbf{a} \cdot \mathbf{b} = 0 .$$

- (viii) \mathbf{a} and \mathbf{b} are parallel if $\theta = 0$, or $\sin \theta = 0$, or

$$\mathbf{a} \times \mathbf{b} = \mathbf{0} .$$

Solution to Exercise 2

- (i) If
- $\mathbf{a} = \mathbf{i} + \mathbf{j}$
- , then
- $|\mathbf{a}| = \sqrt{1^2 + 1^2}$
- , or,

$$|\mathbf{a}| = \boxed{\sqrt{2}}.$$

- (ii)
- $\mathbf{a} - 2\mathbf{b} = -3\mathbf{i} + 3\mathbf{j} + 2\mathbf{k}$
- ,

so that

$$|\mathbf{a} - 2\mathbf{b}| = \sqrt{3^2 + 3^2 + 2^2},$$

or

$$|\mathbf{a} - 2\mathbf{b}| = \boxed{\sqrt{22}}.$$

- (iii)
- $\mathbf{a} \cdot \mathbf{b} = (\mathbf{i} + \mathbf{j}) \cdot (2\mathbf{i} - \mathbf{j} - \mathbf{k}) = 2 - 1$
- ,

so that

$$\mathbf{a} \cdot \mathbf{b} = \boxed{1}.$$

- (iv)
- $\mathbf{a} \times \mathbf{b} = (\mathbf{i} + \mathbf{j}) \times (2\mathbf{i} - \mathbf{j} - \mathbf{k})$
- .

Using the formulae for vector products involving the unit vectors, \mathbf{i} , \mathbf{j} and \mathbf{k} (see page 8 of *SS* if necessary) we see that this is equal to

$$-\mathbf{i} \times \mathbf{j} - \mathbf{i} \times \mathbf{k} + \mathbf{j} \times 2\mathbf{i} - \mathbf{j} \times \mathbf{k} = -\mathbf{k} + \mathbf{j} - 2\mathbf{k} - \mathbf{i},$$

so that

$$\mathbf{a} \times \mathbf{b} = \boxed{-\mathbf{i} + \mathbf{j} - 3\mathbf{k}}.$$

- (v)
- $\mathbf{a} \cdot \mathbf{b} \times \mathbf{c} = (\mathbf{i} + \mathbf{j}) \cdot (2\mathbf{i} - \mathbf{j} - \mathbf{k}) \times (-\mathbf{i} + 3\mathbf{j} + \mathbf{k})$
- .

Proceeding as in part (iv),

$$\begin{aligned} & (2\mathbf{i} - \mathbf{j} - \mathbf{k}) \times (-\mathbf{i} + 3\mathbf{j} + \mathbf{k}) \\ &= 2\mathbf{i} \times 3\mathbf{j} + 2\mathbf{i} \times \mathbf{k} - \mathbf{j} \times (-\mathbf{i}) - \mathbf{j} \times \mathbf{k} - \mathbf{k} \times (-\mathbf{i}) - \mathbf{k} \times 3\mathbf{j} \\ &= 6\mathbf{k} - 2\mathbf{j} - \mathbf{k} - \mathbf{i} + \mathbf{j} + 3\mathbf{i} \\ &= 2\mathbf{i} - \mathbf{j} + 5\mathbf{k}. \end{aligned}$$

Then

$$(\mathbf{i} + \mathbf{j}) \cdot (2\mathbf{i} - \mathbf{j} + 5\mathbf{k}) = 2 - 1,$$

so that

$$\mathbf{a} \cdot \mathbf{b} \times \mathbf{c} = \boxed{1}.$$

Solution to Exercise 3

- (i) The velocity
- \mathbf{v}
- of the particle is equal to
- $d\mathbf{r}/dt$
- . This is

$$\boxed{-6 \sin 3t \mathbf{i} + 6 \cos 3t \mathbf{j} + 2\mathbf{k}}.$$

- (ii) The speed is the magnitude of the velocity. Since

$$|\mathbf{v}| = \sqrt{36 \sin^2 3t + 36 \cos^2 3t + 4},$$

the speed is

$$\boxed{\sqrt{40}}.$$

- (iii) Since the particle is travelling in the direction of its velocity vector at any instant, the unit vector required is $\mathbf{v}/|\mathbf{v}|$. From parts (i) and (ii) this is

$$\frac{1}{\sqrt{40}}(-6 \sin 3t \mathbf{i} + 6 \cos 3t \mathbf{j} + 2 \mathbf{k})$$

0S.4.2 Line Integrals

Solution to Exercise 1

Since

$$\begin{aligned}\mathbf{F} \cdot d\mathbf{r} &= (x\mathbf{i} + x^2y\mathbf{j}) \cdot (\mathbf{i} dx + \mathbf{j} dy) \\ &= x dx + x^2y dy,\end{aligned}$$

and since $dy = 0$ on the line segment from $(-1, 0)$ to $(1, 0)$,

$$\begin{aligned}\int_C \mathbf{F} \cdot d\mathbf{r} &= \int_{-1}^1 x dx \\ &= \left[\frac{1}{2}x^2\right]_{-1}^1,\end{aligned}$$

so that

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \boxed{0}.$$

Solution to Exercise 2

Clearly, from the answer to Exercise 1, the integral is equal to the integral along the line segment from $(1, 0)$ to $(1, 1)$. Since $x = 1$ and $dx = 0$ here,

$$\begin{aligned}\int_C \mathbf{F} \cdot d\mathbf{r} &= \int_0^1 y dy \\ &= \left[\frac{1}{2}y^2\right]_0^1,\end{aligned}$$

so that

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \boxed{\frac{1}{2}}.$$

Solution to Exercise 3

Here we have

$$\frac{dx}{dt} = 2t, \frac{dy}{dt} = 3t^2$$

along C , so that

$$\begin{aligned}\int_C \mathbf{F} \cdot d\mathbf{r} &= \int_C x dx + \int_C x^2y dy \\ &= \int_0^1 t^2 \cdot 2t dt + \int_0^1 t^4 \cdot t^3 \cdot 3t^2 dt \\ &= \int_0^1 (2t^3 + 3t^9) dt \\ &= \left[\frac{1}{2}t^4 + \frac{3}{10}t^{10}\right]_0^1,\end{aligned}$$

so that

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \boxed{\frac{4}{5}}.$$

Solution to Exercise 4

The semicircle is given in Cartesian coordinates by

$$x^2 + y^2 = 1, y \geq 0.$$

We see that $x^2 + y^2$ will be equal to unity if we put

$$x = \cos t, y = \sin t,$$

and that t will vary from 0 to π as the point (x, y) moves around the semicircle from $(1, 0)$ to $(-1, 0)$. Thus our parametric representation of the semicircle is

$$x = \boxed{\cos t},$$

$$y = \boxed{\sin t},$$

$$t \in \boxed{[0, \pi]}.$$

Solution to Exercise 5

Here we have

$$\begin{aligned} \mathbf{F} \cdot d\mathbf{r} &= (\mathbf{i} + y\mathbf{j}) \cdot (\mathbf{i} dx + \mathbf{j} dy) \\ &= dx + y dy, \end{aligned}$$

so that

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C dx + \int_C y dy.$$

On the semicircle,

$$\frac{dx}{dt} = -\sin t, \frac{dy}{dt} = \cos t,$$

so that

$$\begin{aligned} \int_C \mathbf{F} \cdot d\mathbf{r} &= -\int_0^\pi \sin t dt + \int_0^\pi \sin t \cos t dt \\ &= [\cos t - \tfrac{1}{4} \cos 2t]_0^\pi, \end{aligned}$$

or

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \boxed{-2}.$$

Solution to Exercise 6

We have

$$\mathbf{v} \times d\mathbf{r} = (xy dy - y^2 dx)\mathbf{k},$$

so that using the parametrization of Exercise 4 for the semicircle,

$$\begin{aligned} \int_C \mathbf{v} \times d\mathbf{r} &= \left\{ \int_C xy dy - \int_C y^2 dx \right\} \mathbf{k} \\ &= \left\{ \int_0^\pi \cos t \sin t \cos t dt - \int_0^\pi \sin^2 t (-\sin t) dt \right\} \mathbf{k} \\ &= \left\{ \int_0^\pi \sin t dt \right\} \mathbf{k} \\ &= [-\cos t]_0^\pi \mathbf{k}, \end{aligned}$$

so that

$$\int_C \mathbf{v} \times d\mathbf{r} = \boxed{2\mathbf{k}}.$$

Since

$$\int_C \mathbf{v} \times d\mathbf{r} = \mathbf{k} \int_C \mathbf{v} \cdot \mathbf{n} \, dr,$$

it follows that

$$\int_C \mathbf{v} \cdot \mathbf{n} \, dr = \boxed{2}.$$

0S.5 DOUBLE AND VOLUME INTEGRALS

0S.5.1 Double Integrals

Solution to Exercise 1

The region of integration is enclosed between the lines $x = 0$, $x = 1$, $y = 0$ and $y = 2x + 1$. A diagram of this region can be found on page 17 of *Unit MST 282 6*, or on page 11 of *Unit 3*.

Solution to Exercise 2

$$\begin{aligned}\int_0^{2x+1} xy \, dy &= x \int_0^{2x+1} y \, dy \\ &= x \left[\frac{1}{2} y^2 \right]_0^{2x+1},\end{aligned}$$

that is,

$$\int_0^{2x+1} xy \, dy = \boxed{\frac{1}{2}x(2x+1)^2}.$$

Solution to Exercise 3

$$\begin{aligned}\int_0^1 \frac{1}{2}x(2x+1)^2 \, dx &= \frac{1}{2} \int_0^1 (4x^3 + 4x^2 + x) \, dx \\ &= \frac{1}{2} \left[x^4 + \frac{4}{3}x^3 + \frac{1}{2}x^2 \right]_0^1 \\ &= \frac{1}{2} \left[1 + \frac{4}{3} + \frac{1}{2} \right],\end{aligned}$$

so that

$$\int_0^1 \left[\int_0^{2x+1} xy \, dy \right] dx = \boxed{\frac{17}{12}}.$$

Solution to Exercise 4

The “inner” integration is performed with respect to y . Since the ellipse can be written as

$$y = \pm b\sqrt{1 - \frac{x^2}{a^2}},$$

the region enclosed by the ellipse can be described by

$$\begin{aligned}-b\sqrt{1 - \frac{x^2}{a^2}} &\leq y \leq b\sqrt{1 - \frac{x^2}{a^2}}, \\ -a &\leq x \leq a.\end{aligned}$$

The limits (end-points) of the two integrations are then the upper and lower bounds in these two inequalities. Therefore:

$$\begin{aligned}\text{Limits of } y\text{-integration are } &\int_{-b\sqrt{1 - x^2/a^2}}^{b\sqrt{1 - x^2/a^2}}; \\ \text{Limits of } x\text{-integration are } &\int_{-a}^a;\end{aligned}$$

Resulting double integral is

$$\int_{-a}^a \left[\int_{-b\sqrt{1-x^2/a^2}}^{b\sqrt{1-x^2/a^2}} \left(1 - \frac{x^2}{a^2} - \frac{y^2}{b^2} \right) dy \right] dx.$$

Solution to Exercise 5

$$\begin{aligned} \int_{-b\sqrt{1-x^2/a^2}}^{b\sqrt{1-x^2/a^2}} \left(1 - \frac{x^2}{a^2} - \frac{y^2}{b^2} \right) dy \\ &= \left[\left(1 - \frac{x^2}{a^2} \right) y - \frac{y^3}{3b^2} \right]_{-b\sqrt{1-x^2/a^2}}^{b\sqrt{1-x^2/a^2}} \\ &= \left[1 - \frac{x^2}{a^2} - \frac{1}{3} \left(1 - \frac{x^2}{a^2} \right) \right] 2b\sqrt{1 - \frac{x^2}{a^2}}, \end{aligned}$$

so that

Inner integral is $\frac{4}{3}b \left(1 - \frac{x^2}{a^2} \right)^{\frac{3}{2}}.$

We are left with the integral

$$\int_{-a}^a \frac{4}{3}b \left(1 - \frac{x^2}{a^2} \right)^{\frac{3}{2}} dx$$

to evaluate. The substitution to make here is $x = a \sin v$. (You may like to refer to section III.3 of *TI* to see why we choose this particular substitution. We can see that our integral is of the form given at the beginning of that section, with $f(x) = 0$, $h(x) = \frac{4}{3}b(1 - x^2/a^2)$, $ax^2 + bx + c$ replaced by $1 - x^2/a^2$, $g(x) = 1$, $k(x) = 0$. Subsection III.3.2(iii) gives the substitution we are using.) The range $-a \leq x \leq a$ is equivalent to $-\pi/2 \leq v \leq \pi/2$, and $dx/dv = a \cos v$. Thus the integral becomes

$$\frac{4}{3}ba \int_{-\pi/2}^{\pi/2} \cos^4 v \, dv = \frac{1}{2}\pi ab,$$

using the information given in the question. Thus:

Final result is $\frac{1}{2}\pi ab.$

Solution to Exercise 6

Putting $x^2 + y^2 = r^2$ and replacing $dx \, dy$ by $r \, dr \, d\theta$,

$$\iint_D \left(1 - \frac{x^2 + y^2}{a^2} \right) dx \, dy = \iint_D \left(1 - \frac{r^2}{a^2} \right) r \, dr \, d\theta.$$

D is composed of points (r, θ) , where

$$0 \leq r \leq a,$$

$$-\pi \leq \theta \leq \pi.$$

Therefore, the double integral in polar coordinates is

$$\int_0^a \left[\int_{-\pi}^{\pi} \left(1 - \frac{r^2}{a^2} \right) r \, d\theta \right] dr.$$

The inner integral is equal to

$$\left(1 - \frac{r^2}{a^2} \right) r \int_{-\pi}^{\pi} d\theta = 2\pi r \left(1 - \frac{r^2}{a^2} \right).$$

The outer integral is

$$\begin{aligned} 2\pi \int_0^a \left(r - \frac{r^3}{a^2} \right) dr &= 2\pi \left[\frac{1}{2}r^2 - \frac{r^4}{4a^2} \right]_0^a \\ &= 2\pi \left[\frac{1}{2}a^2 - \frac{1}{4}a^2 \right], \end{aligned}$$

so that the value of the integral is

$$\boxed{\frac{1}{2}\pi a^2}.$$

This agrees with the result of Exercise 5 when $b = a$.

0S.5.2 Volume Integrals

Solution to Exercise 1

The interior of the cube consists of points (x, y, z) for which

$$0 \leq x \leq 1,$$

$$0 \leq y \leq 1,$$

$$0 \leq z \leq 1.$$

Thus the triple integral is

$$\boxed{\int_0^1 \left\{ \int_0^1 \left[\int_0^1 (x + y + z) dx \right] dy \right\} dz}.$$

Since

$$\int_0^1 (x + y + z) dx = \left[\frac{1}{2}x^2 + (y + z)x \right]_0^1,$$

the inner integral is

$$\boxed{\frac{1}{2} + y + z}.$$

Since

$$\int_0^1 \left(\frac{1}{2} + y + z \right) dy = \left[\left(\frac{1}{2} + z \right)y + \frac{1}{2}y^2 \right]_0^1,$$

the next integral is

$$\boxed{1 + z}.$$

Since

$$\int_0^1 (1 + z) dz = \left[z + \frac{1}{2}z^2 \right]_0^1,$$

the outer integral is

$$\boxed{\frac{3}{2}}.$$

Solution to Exercise 2

The interior of the prism consists of points (x, y, z) for which

$$0 \leq x \leq 1$$

$$0 \leq y \leq 1 - z$$

$$0 \leq z \leq 1,$$

so that the triple integral is

$$\int_{\boxed{0}}^{\boxed{1}} \left\{ \int_{\boxed{0}}^{\boxed{1-z}} \left[\int_{\boxed{0}}^{\boxed{1}} 3x^2 yz \, dx \right] dy \right\} dz.$$

Solution to Exercise 3

$$\int_0^1 3x^2 yz \, dx = [x^3 yz]_0^1,$$

so that the inner integral is

$$\boxed{yz}.$$

$$\int_0^{1-z} yz \, dy = [\tfrac{1}{2}y^2 z]_0^{1-z},$$

so that the next integral is

$$\boxed{\tfrac{1}{2}z(1-z)^2}.$$

$$\begin{aligned} \tfrac{1}{2} \int_0^1 z(1-z)^2 \, dz &= \tfrac{1}{2} \int_0^1 (z - 2z^2 + z^3) \, dz \\ &= \tfrac{1}{2} [\tfrac{1}{2}z^2 - \tfrac{2}{3}z^3 + \tfrac{1}{4}z^4]_0^1, \end{aligned}$$

so that the outer integral is

$$\boxed{\tfrac{1}{24}}.$$

ISBN 0 335 01257 4